

LITERATUR ZU AMNION

Akle, C.A. ; Adolji, M.; Mc Coll, I.

Permeability of the amniotic membrane and its potential application for transplantation purposes.
Biological Research in Pregnancy 2 (1981), 23-37

Aplin, J.D.; Campell; S.; Foden, L.J.

Adhesion of human amnion epithelial cells to extracellular matrix.
Exp. Cell. Res. 153 (1984), 425-438

Burgos, H.

Angiogenic and growth factors in human amnion-chorion and placenta.
Eur. J. Clin. Invest. 13 (1983), 289-296

Egan, T.J.; O'Driscoll, M.B.; Thakar, D.R.

Human amnion in the management of chronic ulceration of the lower limb: a clinico-pathologic study.
Vasa 3 (1983): 197-203

Göbel, P.; Schubert, W.

Unsere Erfahrung mit der Deckung kindlicher Verbrennungswunden durch Amnion.
Beitr. Orthop. Traumatol. 37 (1990), 495-498

Gruss, J.S.; Dennis, B.C.; Jirsch, D.W.

Human amniotic membrane: a versatile wound dressing.
CMA Journal 118 (1978), 1237-1256

Hampson, V.; Lia, D.; Billet; E.; Kirk; S.

Amniotic membrane collagen content and type distribution in women with preterm premature rupture of the membranes in pregnancy.
Br. J. Obstet. Gynecol. 104 (1997): 1087-1091

Kim, J.C.; Tseng, S.C.G.

Transplantation of preserved human amniotic membrane for surface reconstruction in severely damaged rabbit corneas.
Cornea 14 (1995), 473-484

Kucan, J.O.; Robson, M.C.; Parsons, R.W.

Amniotic membranes as dressings following facial dermabrasion.
Ann. Plast. Surg. 8 (1982): 523-527

Lee, S.H.; Tseng, S.C.G.

Amniotic membrane transplantation for persistent epithelial defects with ulceration.
Am. J. Ophthalmol. 123 (1997), 303-312

Noguchi, Y.; Uchida, Y.; Endo, T.; Ninomiya, H.; Nomura, A.; Sakamoto, T.; Goto, Y.; Haraoka, S.; Shimokama, T.; Watanabe, T.; Hasegawa, S.

The induction of cell differentiation and polarity of tracheal epithelium cultured on the amniotic membrane.
Biochem. Biophys. Res. Commun. 219 (1995), 302-309

Paradowska, E.; Blach-Olzewska, Z.; Sender, J.; Jarosz, W.

Antiviral nonspecific Immunity of Human Placenta at Term: Possible Role of Endogenous Tumor Necrosis Factors and Interferons.
J. Interferon Cytokine Res. 16 (1996), 941-948

Prabhasawat, P.; Tesavibul, N.

Preserved amniotic membrane transplantation for conjunctival surface reconstruction.
Cell and Tissue Banking 2 (2000), 31-39

Ramakrishnan, K.M.; Jayaraman, V.

Management of partial-thickness burn wounds by amniotic membrane: a cost-effective treatment in developing countries.
Burns 23 (1997), 33-36

Robeson, M.C.; Krizek, T.J.

The effect of human amniotic membrane on the bacterial population of infected rat burns.
Am. J. Surg. 177 (1973), 144-149

Sawhney, C.P.

Amniotic membrane as a biological dressing in the management of burns.
Burns 15 (1989), 339-342

Van Herendael, B.J.; Oberti, C.; Brosens; I.

Microanatomy of the human amniotic membranes. A light microscopic, transmission and scanning electron microscopy study.
Am. J. Obstet. Gynaecol. 131 (1978), 872-880

von Versen-Höyneck, Frauke; U. Hesselbarth and D. E. Moeller

Application of sterilised human amnion for reconstruction of the ocular surface.
Cell and Tissue Banking 2003; 00: 1-0

von Versen-Höyneck, Frauke; Carina Syring, S. Bachmann and D. E. Moeller

The influence of different preservation and sterilisation steps on the histological properties of amnion allografts - Light and scanning electron microscopic studies.

Cell and Tissue banking 2004; 00: 1-12

Yokoo, K.; Halaban, R.; Cuono, C.B.

Human amnion cells contain basic fibroblast growth factor and hepatocyte growth factor.
Proc. 24th Am. Burn. Assoc. Mtg. (1992), 31

LITERATUR ZU HAUTALLOGRAFTS

Blome-Eberwein, S.; Jester, A.; Kuentscher, M.; Raff, T.; Germann, G.; Pelzer, M.

Clinical practice of glycerol preserved allograft skin coverage.

Burns 28 (2002), S10-12

Brans, T.A.; Hoekstra, M.J.; Vloemans, A.F.P.M.; Kreis, R.W.

Long-term results of treatment of scalds in children with glycerol-preserved allografts.

Burns 20 (1994), S10-13

Cuono, C.; Langdon, R.; McGuire, J.

Use of cultured epidermal autografts and dermal allografts as skin replacement after burn injury

Lancet 11 (1986), 1123

Cuono, C.B.; Langdon, R.; Birchall, N.; Barttelbort, S.; McGuire, J.

Composite autologous-allogeneic skin replacement: development and clinical application.

Plast. Reconstr. Surg. 80 (1987), 626-637

Germann, G.; Raff, T.

Fremdhauttransplantation bei Schwerbranntten.

Chirurg 66 (1995), 260-270

Gher, M.E.; Williams, J.E.; Vernino, A.R.; Strong, D.M.; Pelleu, G.B.

Evaluation of the immunogenicity of freeze-dried skin allografts in humans.

J. Periodontol. 51 (1980), 571-577

Hickerson, W.L.; Compton, C.; Fletchall, S.; Smith, L.R.

Cultured epidermal autografts and allodermis combination for permanent burn wound coverage.

Burns 20 (1994); S52-5

Izmailow, G.A.; Izmailow, S.G.; Popow, A.N.

Kombinierte Autoallodermoplastik: klinische Begründung ihrer Anwendung.

Z. Dermatologie 182 (1986), 13-17

Kreis, R.W.; Mackie, D.P.; Vloemans, A.W.F.P.; Hermans, R.P.; Hoekstra, M.J.

Widely expanded postage stamp skin grafts using a modified Meek technique in combination with an allograft overlay.

Burns 19 (1993), 142-145

Lamke, L.O.

The influence of different „skin grafts“ on the evaporative water loss from burns.

Scand. J. Plast. Reconstr. Surg. 5 (1971), 82-86

Lin, S.D.; Chai, C.Y.; Lai, C.S.; Chou, C.K.

A mixture of allogeneic and autologous microskin grafting of rabbit skin wounds with Biobrane overlay.

Burns 20 (1994), 30-35

Mac Neil, S.

What role does the extracellular matrix serve in skin grafting and wound healing?

Burns 20 (1994), S67-70

Mackie, D.

Postal survey on the use of glycerol-preserved allografts in clinical practice.

Burns 28 (2002), S40-44

Moerman, E.; Middelkoop, E.; Mackie, D.; Groenevelt, F.

The temporary use of allograft for complicated wounds in plastic surgery.

Burns 28 (2002), S13-15

Peeters, R.; De Caluwe, D.; Neetens, C.; Hubens, A.

Use of glycerolized cadaver skin for the treatment of scalds in children.

Burns 20 (1994); S32-33

Raghow, R.

The role of extracellular matrix in postinflammatory wound healing and fibrosis.

FASEB J. 8 (1994), 823-831

Richters, C.D.; Hoekstra, M.J.; van Baare, J.; du Pont, J.S.; Kamperdijk, E.W.A.

Morphology of glycerol-preserved human cadaver skin.

Burns 22 (1996), 113-116

Shira, R.B.; Carroll, P.B.; Tow, H.D.; Vernino, A.R.

The use of allogeneic freeze-dried skin grafts in the oral environment.

Oral Surg. Oral Med. Oral Path. 37 (1974), 163-175

Sieminski, A.L.; Gooch, K.J.

Biomaterial-microvasculature interactions.

Biomaterials 21 (2000), 2232-2241

Smith, L.T.; Holbrook, K.A.; Byers, P.H.

Structure of the dermal matrix during development and in the adult.

J. Invest. Dermatol. 79 (1982), 93s-104s

Snyder, R.J.; Simonson, D.A.

Cadaveric allograft as adjunct therapy for nonhealing ulcers.
J. Foot Ankle Surg. 38 (1999), 93-101

Vloemans, A.F.; Soesman, A.M.; Suijker, M.; Kreis, R.W.; Middelkoop, E.

A randomised clinical trial comparing hydrocolloid-derived dressing and glycerol preserved allograft skin in the management of partial thickness burns.
Burns 29 (2003); 702-710

Wai, R.T.J.

Use of glycerol-preserved skin in plastic surgery.
Burns 20 (1994); S27-31

Woodley, D.T.; O'Keefe, E.J.; Prunieras, M.

Cutaneous wound healing: a model for cell-matrix interactions.
J. Am. Acad. Dermatol. 12 (1985), 420-433

Wu, J.; Barisoni, D.; Armato, U.

An investigation into the mechanisms by which human dermis does not significantly contribute to the rejection of allo-skin grafts.
Burns 21 (1995), 11-16

LITERATUR ZUR QUALITÄTSSICHERUNG IM TISSUE BANKING

Moening, H.-J.; von Versen, R.

Standards, regulatory and quality systems, within a global context.
In: Radiation in Tissue Banking. Glyn O. Phillips (Ed.), World Scientific, 2000

von Versen, R.

Musculoskeletal tissue banking in europe – Regulations and quality assurance.
Annales Chirurgiae et Gynaecologiae 88 (1999), 215–220

von Versen, R.; Mönig, H.J.

Quality management systems in Tissue Banking.
In: Advances in Tissue Banking, Vol. 4, pp. 1 – 23, GOP Phillips, DM Strong, R von Versen, A Nather (Eds), World Scientific Publishers, 1999

von Versen, R.; Mönig, H.J.; Salai, M.; Bettin, D.

Quality issues in tissue banking: Quality management systems – A review.
Cell and Tissue Banking 1 (2000), 181–192

LITERATUR ZUR SICHERHEIT VON ALLOGENEN GEWEBETRANSPLANTATEN

Pruss, A.; Baumann, B.; Seibold, M.; Kao, M.; Tintelnot, K.; von Versen, R.; Radtke, H.; Dörner, T.; Pauli, G.; Göbel, B.

Validation of the sterilization procedure of allogeneic avital bone transplants using peracetic acid-ethanol.
Biologicals 29 (2000), 59–66

Pruss, A.; Göbel, U.B.; Kao, M.; Seibold, M.; Mönig, H.J.; Hansen, A.; von Versen, R.

Peracetic acid-ethanol treatment of allogeneic avital bone tissue transplants – a reliable sterilization method.
Annals of Transplantation 8 (2003), 34–42

Pruss, A.; Kao, M.; Gohs, U.; Koscielny, J.; von Versen, R.; Pauli, G.

Effect of gamma irradiation on human cortical bone transplants contaminated with enveloped and non-enveloped viruses.
Biologicals 30 (2002), 125–133

Pruss, A.; Kao, M.; Kiesewetter, H.; Versen, R.; Pauli, G.

Virus safety of avital bone tissue transplants: Evaluation of sterilization steps of spongiosa cuboids using a peracetic acid-methanol mixture.
Biologicals 27 (1999), 195–201

Pruss, A.; Perka, C.; Degenhardt, P.; Maronna, U.; Buettner-Jan, K.; Pau, B.; Mueller, K.; Klumpp, C.; Bruck, J.C.; von Versen, R.

Clinical efficacy and compatibility of allogeneic avital tissue transplants sterilized with a peracetic acid/ethanol mixture.
Cell and Tissue Banking 3 (2002), 235–243

Salai, M.; Vonsover, A.; Pritch, M.; von Versen, R.; Horoszowski, H.

Human Immunodeficiency virus (HIV) inactivation of banked bone by gamma irradiation.
Annals of Transplantation 2 (1996), 55–56

LITERATUR ZU KERATINOZYTEN

Auböck, J.; Romani, N.; Kompatscher, P.; Sifter, M.; Bauer, M.; Fritsch, P.

Keratinocytenexpansionskultur zur Deckung von Hautdefekten.

In: Fortschritte der praktischen Dermatologie und Venerologie Band, 11: 418 - 427.
Springer-Verlag, Berlin-Heidelberg-New York 1986

Barandon, Y.; Green, H.

Cell migration is essential for sustained growth of keratinocyte colonies: the roles of transforming growth factors-alpha and epidermal growth factor.

Cell 50 (1987), 1131-1137

Ben-Bassat, H.; Eldad, A.; Chaouat, M.; Livoff, A.; Ron, N.; Neeman, Z.; Wexler, M.R.

Structural and functional evaluation of modifications in the composite skin graft: cryopreserved dermis and cultured keratinocytes.

Plast. Reconstr. Surg. 89 (1991), 510-520

Bloemkolk, J.W.; Gray, R.G.

Effect of temperature of hybridoma cycle and mAb production.

Biotechnol. Bioeng. 40 (1992), 427-431

Bödeker, B.G.D.; Steiner, U.; Berg, G.J.; Hewlett, G.; Schlumberger, H.D.

Produktion natürlicher Substanzen aus tierischen Zellkulturen in technischen Kultursystemen.

Tierische Zellkulturen, Bericht vom 2. BMBF-Statusseminar 1985, 7-129

Boisseau, A.M.; Donatien, P.; Surleve-Bazeille, J.E.; Amedee, J.; Harmand, M.F.; Beziau, J.H.; Maleville, J.; Taieb, A.

Production of epidermal sheets in a serum free culture system: a further appraisal of the role of extracellular calcium.

J. Dermatol. Sci. 3 (1992), 111-120

Boyce, S.T.; Ham, R.G.

Calcium-regulated differentiation of normal human epidermal keratinocytes in chemically defined clonal culture and serum-free serial culture.

J. Invest. Dermatol. 81 (1983), 33s-40s

Boyce, S.T.; Kagan, R.J.; Meyer, N.A.; Yakuboff, K.P.; Warden, G.D.

Cultured skin substitutes combined with Integra Artificial Skin to replace native skin autograft and allograft for the closure of excised full-thickness burns.

J. Burn Care Rehabil. 20 (1999), 453-461

Braye, F.; Oddou, L.; Bertin-Maghit, M.

Widely Meshed Autograft Associated with Cultured Autologous Epithelium for the Treatment of Major Burns in Children: Report of 12 Cases.

Eur. J. Pediatr. Surg. 10 (2000), 35-40

Burke, J.F.; Yannas, J.V.; Quinby, W.; Bondoc, C.C.; Jung, W.J.

Successful use of a physiologically acceptable artificial skin in the treatment of extensive burns.

Ann. Surg. 94 (1981), 413-428

Carter, D. M.; Lin, A. N.; Varghese, M. C.

Treatment of junctional epidermolysis bullosa with epidermal autografts.

J. Am. Acad. Dermatol. 172 (1987), 246-250

Carver, N.; Leigh, I. M.

Keratinocyte grafts and skin equivalents.

Int. J. Dermatol. 30 (1991), 540-551

Chan, E.S.; Lam, P.K.; Liew, C.T.; Lau, H.C.; Yen, R.S.; King, W.W.

A new technique to resurface wounds with composite biocompatible epidermal grafts and artificial skin.

J. Trauma 50, (2001): 358-362

Chen, M.; Kasahara, N.; Keene, D.R.; Chan, L.; Hoeffler, W.K.; Finlay, D.; Barcova, M.; Cannon, P.M.; Mazurek, C.; Woodely, D.T.

Restoration of type VII collagen expression and function in dystrophic epidermolysis bullosa.

Nat. Genet. 32 (2002), 670-675

Compton, C. C.

Current concepts in pediatric burn care: the biology of cultured epithelial autografts: an eight-year study in pediatric burn patients.

Eur. J. Pediatr. Surg. 2 (1992) 216-222

Coulomb, B.; Friteau, L.; Baruch, J.; Guilbaud, J.; Chretien-Marquet, B.; Glicenstein, J.; Lebreton-Decoster, C.; Bell, E.; Dubertret, L.

Advantage of the presence of living dermal fibroblasts within in vitro reconstructed skin for grafting in humans.

Plast. Reconstr. Surg. 101 (1998), 1891-1903

Cuono, C. B.; Langdon, R.; Birchall, N.

Composite autologous-allogenic skin replacement: Development and clinical application.

J. Invest. Dermatol. 80 (1987), 626-635

Currie, L.J.; Martin, R.; Sharpe, J.R.; James, S.E.

A comparison of keratinocyte cell sprays with and without fibrin glue.

Burns 29 (2003), 677-685

Danes, B.S.; Broadfoot, M.M.; Paul, J.

A comparative study of respiratory metabolism in cultured mammalian cell strains.

Exp. Cell. Res. 30 (1963), 369-378

Dantzer, E.; Braye, F.M.

Reconstructive surgery using an artificial dermis (Integra): results with 39 grafts.
Br. J. Plast. Surg. 54 (2001), 659-664

Dawson, R.A.; Goberdhan, N.J.; Freedlander, E.; MacNeil, S.

Influence of extracellular matrix proteins on human keratinocyte attachment, proliferation and transfer to a dermal wound model.
Burn 22 (1996), 93-100

De Luca, M.; Cancedda, R.

Culture of human epithelium.
Burns. 18 (1992); S5-10

Doebis, C.; Ritter, T.; Brandt, C.; Schonberger, B.; Volk, H.D.; Seifert, M.

Efficient in vitro transduction of epithelial cells and keratinocytes with improved adenoviral gene transfer for the application in skin tissue engineering.
Transpl. Immunol. 9 (2002), 323-329

Donati, L.; Magliacani, G.; Bormioli, M.; Signorini, M.; Baruffaldi Preis, F. W.

Clinical experiences with keratinocyte grafts.
Burns. 18 (1992), S19-25

Dvorankova, B.; Holikova, Z.; Vacik, J.; Konigova, R.; Kapounkova, Z.; Michalek, J.; Pradn, M.; Smetana, K.

Reconstruction of epidermis by grafting of keratinocytes cultured on polymer support-clinical study.
Int. J. Dermatol. 42 (2003), 219-223

Eagle, H.

The effect of environmental pH on the growth of normal and malignant cells.
J. Cell. Physiol. 82 (1973), 1-8

Erdag, G.; Medalie, D.A.; Rakhorst, H.; Krueger, G.G.; Morgan, J.R.

FGF-7 expression enhances the performance of bioengineered skin.
Mol. Ther. 10 (2004), 76-85

Frame, J.D.; Still, J.; Lakhel-LeCoadou, A.; Carstens, M.H.; Lorenz, C.; Orlet, H.; Spence, R.; Berger, A.C.; Dantzer, E.; Burd, A.

Use of dermal regeneration template in contraction release procedures: a multicenter evaluation.
Plast. Reconstr. Surg. 113 82004), 1330-1338

Freshney, R.I.

Culture of Animal Cells.
John Wiley & Sons, N.Y.; Chichester, Brisbane, Toronto, Singapore, Third Edition 1994

Galassi, G.; Burn, P.; Radice, M.; Cortivo, R.; Zanon, G.F.; Genovese, P.; Abatangelo, G.

In vitro reconstructed dermis implanted in human wounds: degradation studies of the HA-based supporting scaffold.
Biomaterials 21 (2000), 2183-2191

Gallico, G. G.; O'Connor, N. E.; Compton, C. C.

Cultured epithelial autografts for giant congenital naevi.
Plast. Reconstr. Surg. 9 (1989), 1-9

Guo, M.; Toda, K.I.; Grinnell, F.

Activation of human keratinocyte migration on type I collagen and fibronectin.
J. Cell Sci. 96 (1990), 197-205

Gustavson, C.J.; Kratz, G.

Cultured autologous keratinocytes on a cell-free dermis in the treatment of full-thickness wounds.
Burns 25 (1999), 331-335

Green, H.; Kehinde, O.; Thomas, J.

Growth of cultured human epidermal cells into multiple epithelia suitable for grafting.
Proc. Natl. Acad. Sci. USA 76 (1979), 5665-5668

Haas, A. F., Reilly, D. A.

Cultured epithelial autografts in the treatment of extensive recalcitrant keloids.
Arch. Dermatol. 134 (1998), 549-552

Harriger, M.D.; Supp, A.P.; Warden, G.D.; Boyce, S.T.

Glutaraldehyde crosslinking of collagen substrates inhibits degradation in skin substitutes grafted to athymic mice.
J. Biomed. Mater. Res. 35 (1997), 137-145

Hefton, J. M., Caldwell, D., Biozes, D. G.

Grafting of skin ulcers with cultured autologous epidermal cells.
J. Am. Acad. Dermatol. 14 (1986), 399-405

Henckel von Donnersmark, G.; Mühlbauer, W.; Höfer, E.; Hartinger, A.

Die Verwendung von Keratinozytenkulturen in der Schwerbrandverletztenbehandlung – bisherige Erfahrungen, Ausblicke zur weiteren Entwicklung.
Unfallchirurg 98 (1995), 229-232

Herson, M.R.; Mathor, M.B.; Altran, S.; Capelozzi, V.L.; Ferreira, M.C.

In vitro construction of a potential skin substitute through direct human keratinocyte plating into decellularized glycerol-preserved allodermis.
Artif. Org. 25 (2001), 901-906

Hickerson, W. L.; Compton, C.; Fletchall, Smith, L.R.

Cultured epidermal autografts and allodermis combination for permanent burn wound coverage.
Burns. 20 (1994), S52-55

Holzmann, H.; Kippenberger, S.; Ramirez-Bosca, A.; Bereiter-Hahn, J.; Bernd, A.

Zell- und Gewebekulturmodelle in der Dermatologie.
Hautarzt 45 (1994), 304-312

Hunyadi, J.; Farkas, B.; Bertenyi, C.

Keratinocyte grafting: A new means of transplantation for full thickness wounds.
J. Dermatol. Surg. Oncol. 14 (1988), 75-78

Jones, I.; James, S.E.; Rubin, P.; Martin, R.

Upward migration of cultured autologous keratinocytes in Integra artificial skin: a preliminary report.
Wound Repair Regen. 11 (2003), 132-138

Kopp, J.; Jeschke, M.G.; Bach, A.D.; Kneser, U.; Horch, R.E.

Applied tissue engineering in the closure of severe burns and chronic wounds using cultured human autologous keratinocytes in a natural fibrin matrix.
Cell Tissue Bank. 5 (2004), 89-96

Kraut, J.D.; Eckhardt, A.J.; Patton, A.J.; Antoniadis, K.; Haith, L.R.; Shotwell, B.S.

Combined simultaneous application of cultured epithelial autografts and AlloDerm®.
Wounds 7 (1995), 137-142

Krejci, N.C.; Cuono, C.B.; Langdon, R.C.; McGuire, J.

In vitro reconstitution of skin: fibroblasts facilitate keratinocyte growth and differentiation on acellular reticular dermis.
J. Invest. Dermatol. 97 (1991), 843-848

Kremer, M.; Lang, E.; Berger, A.

Organotypical engineering of differentiated composite-skin equivalents of human keratinocytes in a collagen-GAG matrix (INTEGRA Artificial skin) in a perfusion culture system.
Langenbecks Arch. Surg. 386 (2001), 357-363

Krupp, S.; Benathan, M.; Meuli, M.

Current concepts in pediatric burn care: management of burn wounds with cultured epidermal autografts.
Eur. J. Pediatr. Surg. 2 (1992), 210-215

Leigh, I. M.; Purkis, P. E.

Culture grafted leg ulcers.
Clin. Exp. Dermatol. 11 (1986), 650-652

Llames, S.G.; Del Rio, M.; Larcher, F.; Garcia, E.; Garcia, M.; Escamez, M.J.; Jorcano, J.L.; Holguin, P.; Meana, A.

Human plasma as a dermal scaffold for the generation of a completely autologous bioengineered skin.
Transplantation 77 (2004), 350-5

Lidl, T.

Zell- und Gewebekultur.
Gustav Fischer Verlag, Stuttgart, Jena, New York, 3. Auflage, 1994

Limat, A.; French, L.E.; Blal, L.; Suarat, J.H.; Hunziker, T.; Salomon, D.

Organotypic cultures of autologous hair follicle keratinocytes for the treatment of recurrent leg ulcers.
J. Am. Acad. Dermatol. 48 (2003), 207-214

Lin, A. N.; Balin, A. K.; Pratt, L.; Eisinger, M.; Caldwell, D.; Marko, O.; Carter, D.M.

Treatment of junctional epidermolysis bullosa with epidermal autografts.
Clin. Res. 34 (1986), 764A

Liu, J.Y.; Hafner, J.; Dragieva, G.; Seifert, B.; Burg, G.

Autologous keratinocytes on porcine gelatin microbeads effectively heal chronic leg ulcers.
Wound Repair Regen. 12 (2004), 148-156

Livesey, S.A.; Herndon, D.N.; Hollyoak, M.A.; Atkinson, Y.H.; Nag, A.

Transplanted acellular allograft dermal matrix.
Transplantation 60 (1995), 1-9

Mass-Szabowski, N.; Zabowski, A.; Stark, H.J.; Andrecht, S.; Kolbus, A.; Schorpp-Kistner, M.; Angel, P.; Fusenig, N.

Organotypic cocultures with genetically modified mouse fibroblasts as a tool to dissect molecular mechanisms regulating keratinocyte growth and differentiation.
J. Invest. Dermatol. 116 (2001), 816-820

McAree, K. G.; Klein, R. L.; Boeckman, C. R.

The Use of Cultured Epithelial Autografts in the Wound Care of Severely Burned Patients.
J. Pediatric Surg. 28 (1993), 166-168

Medalie, D.A.; Eming, S.A.; Collins, M.E.; Tompkins, R.G.; Yarmush, M.L.; Morgan, J.R.

Differences in dermal analogs influence subsequent pigmentation, epidermal differentiation, basement membrane, and rete ridge formation of transplanted composite skin grafts.
Transplantation 64 (1997), 454-465

Meng, X.; Sawamura, D.; Ina, S.; Tamai, K.; Hanada, K.; Hashimoto, I.

Keratinocyte gene therapy; cytokine gene expression in local keratinocytes and in circulation by introducing cytokine genes into skin.
Exp. Dermatol. 11 (2002), 456-461

Miller, W.M.; Blanch, H.W.; Wilke, C.R.

A kinetic analysis of hybridoma growth and metabolism in batch and continuous suspension culture: effect of nutrient concentration, dilution rate, and pH.
Biotechnol. Bioeng. 32 (1989 b), 947-965

Miller, W.M.; Wilke, C.R.; Blanch, H.W.

Effects of dissolved oxygen concentration on hybridoma growth and metabolism in continuous culture.
J. Cell. Physiol. 132 (1989 a), 524-530

Moll, I.; Schönfeld, M.; Jung, E. G.

Applikation von Keratinozyten in der Therapie von Ulcera crurum.
Hautarzt 46 (1995), 548-552

Munster, A. A.

Cultured Skin for Massive Burns - A Prospective, Controlled Trial.
Ann. Surg. 224 (1997), 372-377

Navsaria, H. A.; Myers, S. R.; Leigh, I. M.; McKay, I. A.

Culturing skin in vitro for wound therapy.
TIBTECH 13 (1995), 91-100

O'Connor, N. E.; Mulliken, J. B.; Banks-Schlegel, S.

Grafting of burns with cultured epithelium prepared from autologous epidermal cells.
Lancet. 1 (1981), 75-78

Ojeh, N.O.; Frame, J.D.; Navsaria, H.A.

In vitro characterization of an artificial dermal scaffold.
Tissue Engineering 7 (2001), 457-472

Ortiz-Urda, S.; Lin, Q.; Yant, S.R.; Keene, D.; Kay, M.A.; Khavari, P.A.

Sustainable correction of junction epidermolysis bullosa via transposon-mediated nonviral gene transfer.
Gene Ther. 10 (2003), 1099-1104

Pandya, A.N.; Woodward, B.; Parkhouse, N.

The use of cultured autologous keratinocytes with integra in the resurfacing of acute burns.
Plast. Reconstr. Surg. 102 (1998), 825-888

Pillai, S.; Bikle, D.D.; Hincenbergs, M.; Elias, P.M.

Biochemical and morphological characterization of growth and differentiation of normal human neonatal keratinocytes in a serum-free medium.
J. Cell. Physiol. 134 (1988), 229-237

Pittelkow, M. R.; Scott, R. E.

New techniques for the in vitro culture of human skin keratinocytes and perspectives on their use for grafting of patients with extensive burns.
Mayo Clin. Proc. 61 (1986), 771-777

Premachandra, D. J.; Woodward, B. M.; Milton, C. M.

Treatment of postoperative otorrea by grafting of mastoid cavities with cultured autologous epidermal cells.
Lancet. 335 (1990), 365-367

Prenosil, J.E.; Kinooka, M.

Computer controlled bioreactor for large-scale production of cultured skin grafts.
Ann. NY Acad. Sci. 875 (1999), 386-397

Prunieras, M.; Regnier, M.

Epidermal keratinocytes as experimental models.
Models Dermatol. 3 (1987), 8-15

Rennekampff, H.O.; Kiessig, V.; Griffey, S.; Greenleaf, J.; Hansbrough, F.

Acellular dermis promotes cultured keratinocytes engraftment.
J. Burn Care Rehabil. 18 (1997), 535-544

Rennekampff, H.O.; Kiessig, V.; Griffey, S.; Greenleaf, G.; Hansbrough, J.F.

Acellular human dermis promotes cultured keratinocyte engraftment.
J. Burn Care Rehabil. 18 (1997), 535-44

Rheinwald, J.G.

Human epidermal keratinocyte cell culture and xenograft systems: applications in the detection of potential chemical carcinogens and the study of epidermal transformation.
Skin carcinogenesis: mechanisms and human relevance, 113-125, 1989, Alan R. Liss, Inc.

Rheinwald, J. G.; Green, H.

Serial cultivation of strains of human epidermal keratinocytes: The formation of keratinising colonies from single cells.
Cell 6 (1975), 331-344

Rosdy, M.; Clauss, L.C.

Terminal epidermal differentiation of human keratinocytes grown in chemically defined medium on inert filter substrates at the air liquid interface.
J. Invest. Dermatol. 95 (1990), 409-414

Schattenkirchner, S.; Eming, S.; Hunzelmann, N.; Krieg, T.; Smola, H.

Treatment of epidermolysis bullosa acquisita with mycophenolate mofetil and autologous keratinocyte grafting.
Br. J. Dermatol. 141 (1999), 932-933

Schechner, J.S.; Crane, S.K.; Wang, F.; Szeglin, A.M.; Tellides, G.; Lorber, M.I.; Bothwell, A.L.; Pober, J.S.
Engraftment of a vascularized human skin equivalent.
FASEB J. 17 (2003), 2250-2256

Sebök, B.; Török, Z.; Schneider, I.
Chirurgische Entfernung von Tätowierungen, Verbesserung der Ergebnisse mittels Zellsuspensions-Wundbehandlung.
Akt. Dermatol. 16 (1990), 295-297

Sheridan, R.L.; Hegarty, M.; Tompkins, R.G.; Burke, J.F.
Artificial skin in massive burns-results to ten years.
Eur. J. Plast. Surg. 17 (1994), 91-93

Sheridan, R. L.; Tompkins, R. G.
Skin Substitutes in Burns.
Burns 25 (1999), 97-103

Shiple, G.D.; Pittelkow, M.
Control of growth and differentiation in vitro of human keratinocytes cultured in serum-free medium.
Arch. Dermatol. 123 (1987), 1541a-1544a

Somers, T.; Verbeken, G.; Vanhalle, S.; Delaey, B.; Duinslaeger, L.; Govaerts, P.; Offeciers, E.
Treatment of chronic postoperative otorrhea with cultured keratinocyte sheets.
Ann. Otol. Rhinol. Laryngol. 106 (1997), 15-21

Stern, R.; McPherson, M.; Longaker, M.T.
Histologic study of artificial skin used in the treatment of full-thickness thermal injury.
J. Burn Care Rehabil. 11 (1990), 7-13

Still, J. M.; Orlet, H. K.; Law, E. J.
Use of cultured epidermal autografts in the treatment of large burns.
Burns 20 (1994), 539-541

Sugita, T.; Huang, C.C.; Abramson, M.
The effect of fibronectin on the migration of keratinocytes.
Acta Otolaryngol. (Stockh.) 104 (1987), 521-525

Supp, D.M.; Wilson-Landy, K.; Boyce, S.T.
Human dermal microvascular endothelial cells from vascular analogs in cultured skin substitutes after grafting to athymic mice.
FASEB J. 16 (2002); 797-804

Tausche, A.K.; Skaria, M.; Bohlen, L.; Liebold, K.; Hafner, J.; Friedlein, H.; Meurer, M.; Goedkoop, R.J.; Wollina, U.; Salomon, D.; Hunziker, T.
An autologous epidermal equivalent tissue-engineered from follicular outer root sheath keratinocytes is as effective as split-thickness skin autograft in recalcitrant vascular leg ulcers.
Wound Repair Regen. 11 (2003), 248-252

Teepe, R. G. C.; Koebrugge, E. J.; Löwik, C. W. G. M.
Cytotoxic effects of topical antimicrobial and antiseptic agents on human keratinocytes in vitro.
J. Trauma 35 (1993): 8-19

van den Bogaardt, A.J.; Ulrich, M.M.; van Galen, M.J.; Reijnen, L.; Verkerk, M.; Pieper, J.; Lamme, E.N.; Middelkoop, E.
Upside-down transfer of porcine keratinocytes from a porous, synthetic dressing to experimental full-thickness wounds.
Wound Repair Regen. 12 (2004), 225-234

Vogt, P.M.; Hebebrand, D.; Steinau, H.U.
Biologische und molekularbiologische Aspekte der Verbrennungstherapie
Chirurg 66 (1995), 251-259

Voigt, M.; Andree, C.; Kalt, T.; Dormann, S.; Schaefer, D.J.; Walgenbach, K.J.; Stark, G.B.
Human recombinant EGF protein delivered by a biodegradable cell transplantation system.
Tissue Eng. 8 (2002); 263-272

Walter, R.J.; Matsuda, T.; Reyes, H.M.; Walter, J.M.; Hanumadass, M.
Characterization of acellular dermal matrixes (ADMs) prepared by two different methods.
Burns 24 (1998), 104-113

Wille, J.J.; Pittelkow, M.R.; Shiple, G.D.; Scott, R.E.
Integrated control of growth and differentiation of normal human prokeratinocytes cultured in serum-free medium: clonal analysis, growth kinetics, and cell cycle studies.
J. Cell. Physiol. 121 (1984), 31-44

Wisser, D.; Steffes, J.
Skin replacement with a collagen based dermal substitute, autologous keratinocytes and fibroblasts in burn trauma.
Burns 29 (2003), 375-380

Wright, K.A.; Nadire, K.B.; Busto, P.; Tubo, R.; McPherson, J.M.; Wentworth, B.M.
Alternative delivery of keratinocytes using a polyurethane membrane and the implications for its use in the treatment of full-thickness burn injury.
Burns 24 (1998), 7-17

Yannas, I.V.; Lee, E.; Orgill, D.P.; Skrabut, E.M.; Murphy, G.F.
Synthesis and characterization of a model extracellular matrix that induces partial regeneration of adult mammalian skin.
Proc. Natl. Acad. Sci. USA. 86 (1989), 933-937